



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON THE CONVERGENCE OF DEFINITE INTEGRALS.

BY J. K. WHITTEMORE.

LET $f(x)$ be a function of x continuous for all values of x greater than a . Then a sufficient condition for the convergence of the improper definite integral,

$$\int_b^{\infty} f(x) \, dx \quad b > a > 0$$

is

$$(1) \quad \lim_{x=\infty} \left(x^k f(x) \right) = A \quad k > 1$$

where A is any finite number. If this condition is satisfied, then

$$(2) \quad \lim_{x=\infty} \left(x f(x) \right) = 0.$$

Equation (2) is, however, not a sufficient condition. For it is satisfied by $f(x) = \frac{1}{x \log x}$, and the integral

$$\int_2^{\infty} \frac{dx}{x \log x}$$

is divergent. Moreover equation (2) is not a necessary condition, for if

$f(x) = \frac{\sin x}{x}$, the condition (2) is not satisfied; but the integral

$$\int_0^{\infty} \frac{\sin x}{x} \, dx = \frac{\pi}{2}.$$

Equation (2) is not a necessary condition for the convergence of the integral even when the integrand does not change sign. For the integral

$$\int_0^{\infty} \frac{dx}{1+x^4 \sin^2 x}$$

is convergent,* and the expression, $\frac{x}{1+x^4 \sin^2 x}$, approaches no limit as x is indefinitely increased. Equation (2) is, however, a necessary condition for the convergence of the integral:

$$\int_b^{\infty} f(x) dx$$

if $xf(x)$ has not an infinite number of maxima and minima for $x > b$. It is easy to see that in the two examples:

$$xf(x) = \sin x, \quad xf(x) = \frac{x}{1+x^4 \sin^2 x}$$

the function has an infinite number of maxima and minima beyond every value of x . The assumptions we make are the following:

1. $f(x)$ is continuous for $x > a$.
 2. $\int_b^{\infty} f(x) dx$ is convergent, and
 3. $xf(x)$ has not an infinite number of maxima and minima for $x > b > a$.
- Then, we wish to prove that:

$$\lim_{x=\infty} (xf(x)) = 0.$$

A necessary and sufficient condition for the convergence of the integral,

$$\int_b^{\infty} f(x) dx$$

is

$$(3) \quad \lim_{\substack{a_1 = \infty \\ a_2 = \infty}} \left(\int_{a_1}^{a_2} f(x) dx \right) = 0 \quad b < a_1 < a_2$$

* See Appell — *Eléments d'Analyse mathématique*, p. 243.

From (3) it follows that to every positive ϵ there corresponds a quantity $d > b$ such that for $d < a_1 < a_2$,

$$\left| \int_{a_1}^{a_2} f(x) dx \right| < \epsilon.$$

Since $1/x$ does not change sign between $x = a_1$ and $x = a_2$ we may apply the mean value theorem for integrals. Then

$$\begin{aligned} \int_{a_1}^{a_2} f(x) dx &= \int_{a_1}^{a_2} x f(x) \frac{dx}{x} \\ &= \xi f(\xi) \int_{a_1}^{a_2} \frac{dx}{x} & a_1 < \xi < a_2 \\ &= \xi f(\xi) \log \frac{a_2}{a_1}. \end{aligned}$$

Let us choose $a_1 = 2d$, $a_2 = 2e d$, where e is the exponential base.

Then we have, as required, $d < a_1 < a_2$, and $\log \frac{a_2}{a_1} = 1$, and

$$(4) \quad \left| \int_{a_1}^{a_2} f(x) dx \right| = \left| \xi f(\xi) \right| < \epsilon.$$

We may now choose d so large that $xf(x)$ has neither maximum nor minimum for $x > d$, since by hypothesis there are not an infinite number of such points. Then $|xf(x)|$ must either never increase or never decrease as x increases indefinitely, and from (4) it follows that it must decrease indefinitely:

$$\lim_{x = \infty} (xf(x)) = 0.$$

In the same way we may see that a necessary condition for the convergence of the integral,

$$\int_a^b f(x) dx$$

where $f(x)$ is continuous for all values of x satisfying the inequalities,

$$a < x \leq b,$$

and where $f(a) = \infty$, if $(x - a) f(x)$ has not an infinite number of maxima and minima in the neighborhood of $x = a$, is

$$\lim_{x = a} \left[(x - a) f(x) \right] = 0.$$

HARVARD UNIVERSITY, JUNE, 1900.